



# International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR)

[Impact Factor – 0.852]

Journal Homepage: [www.eijppr.com](http://www.eijppr.com)

## Research Article

# Effect of Aluminium Chloride and Protective Effect of Ginger Extract on Hematological Profiles in Male Wistar Rats

Kalaiselvi A<sup>1,\*</sup>, Aadhinaath Reddy G<sup>2</sup>, Ramalingam V<sup>1</sup>

<sup>1</sup>Department of Zoology, K.M.Centre for Post Graduate Studies, Pudukchery- 605008, India

<sup>2</sup>Department of Pharmacology, Siddha Central Research Institute, Chennai, Tamilnadu, India

### Article info

#### Article History:

Received 12 September 2014

Accepted 14 January 2015

#### Keywords:

Aluminium chloride, Ginger, Rat, RBC, WBC, Hemoglobin, Hematocrit

### Abstract

The present study was planned to investigate the effect of aluminium chloride and the protective effect of ginger extract on hematological profiles in male rats. One group of rats was given aluminium chloride orally (100mg/kg bw) for 60 days, another group with aluminium chloride and ginger extract (50mg/kg bw) simultaneously and the third group with ginger extract alone. All the three groups were compared with one control group. The hematological parameters were: total erythrocytes count (RBC), White blood cells (WBC), hemoglobin(HB), hematocrit (HCT), and red blood indices: mean corpuscular volume (MCV) mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHV) was investigated. The obtained results showed significant increase in WBC count and decrease in HCT, HB and RBC. No change was noted in (MCV) and there was reduction in mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHV) in AlCl<sub>3</sub> administered group (E1). However, no appreciable changes were noted in AlCl<sub>3</sub> and ginger extract treated (E2) and Ginger alone treated group (E3). The changes in the values of blood indices may be a defensive mechanism against aluminum toxicity through stimulation of erythropoiesis. It may be concluded that Ginger rich in antioxidant and also spicy content which is safety food for human consumption. This study showed that hematologic effects of AlCl<sub>3</sub> are reversible by aqueous ginger extract.

## 1. INTRODUCTION

The increase in pollution is a major global problem and this is due to the use of toxic chemicals or xenobiotic substances or by certain synthetic compounds such as heavy metallic compounds<sup>1,2</sup>. Aluminum (Al) is the third most prevalent and the most abundant metal in the earth's crust, representing approximately 8% of total mineral components<sup>3</sup>. Al components are primarily found in corn, yellow cheese, salt, herbs, spices, tea. It is widely used in cosmetics, cookware utensils and containers, food additives and toothpaste<sup>4</sup> and used in medicines, e.g., antacids, phosphate binders, buffered aspirins, vaccines and injectable allergens<sup>5,7</sup> and also added to drinking water for purification purposes<sup>8</sup>. Recently, increased attention is being paid to aluminum due to its serious effects on the central nervous system, energy metabolism and hematology. The different forms of Al have been shown to be systemic toxicants<sup>9</sup> and there is now considerable evidence demonstrating that the accumulation of Al in body tissues is associated with damage to the target organs<sup>10</sup>.

Blood, which rapidly and constantly flows through the brain and other body organs and play an important role in the transportation of nutrients, antioxidants, hormones and some other chemicals. Once in the blood circulation, Al is mainly transported by plasma transferrin in its sites left vacant by iron, and to a much lesser extent by albumin<sup>11</sup>. Hematological and biochemical profiles of blood can provide important information about the internal environment of the organism<sup>12</sup>. Hematological parameters as red blood cells counting (RBC), hemoglobin concentration (HB), hematocrit(HCT), leukocytes, neutrophils and mononuclear cells counting and Mean Corpuscular Volume (MCV) are largely applied to diseases diagnostics, including effects due to exposure to toxic substances.

Now a day's lot of research has been conducted on the use of herbal products as natural antioxidants because of their fewer side effects, easy and cheap availability. Medicinal plants have been used in various traditional systems, as they have immune potential against numerous diseases<sup>13</sup>. Ginger (*Zingiber officinale*) is commonly used as food spice since in ancient times<sup>14</sup>. It has long been used as a remedy for common ailments like digestive problems, cold, fever, morning sickness. All major active ingredients of ginger such as zingerone, gingerdiol, zingibrene, gingerols and shogaols have antioxidant activity<sup>15</sup>. Studies have revealed a wide range of pharmacological effects demonstrated are anti-platelet, antioxidant, anti-tumour, anti-rhinoviral, anti-hepatotoxicity and anti-arthritis effect<sup>16-20</sup>.

Ginger can be successfully used as a component of curry powder, sauces, ginger bread and ginger flavoured carbonated drinks and in preparation of dietaries for its aroma and flavor. With regard to antioxidant properties of ginger, it can be selected for this study to evaluate the ameliorative effects. Therefore, the present study was carried out to investigate the effect of aluminium chloride in blood hematological profile and the possible protective effect of ginger extracts on it.

## 2. MATERIALS AND METHODS

### 2.1 Animals

Adult albino rats (Wistar strain) weighing 250-260g were used in this study. Animals were maintained as per the guidelines of the Institutional Animal Ethical Committee (IAEC). All the animals were housed in polypropylene cages at standard husbandry conditions (Temperature: 23 ± 2°C, Relative humidity: 60-70%, 12h: 12h light /dark cycle) and were provided with standard pellets and water ad libitum. They were initially acclimatized for the study. The experimental protocols were approved by the Institutional Animal Ethical Committee (IAEC) Approval No 140/PHARMA/SCRI, 2013. After 15 days adaptation period the rats were divided into four groups of 6 animals each.

#### \*Corresponding Author:

Department of Zoology,  
K.M.Centre for Post Graduate Studies,  
Pudukchery- 605008, India.

Email: [pravinakalai@gmail.com](mailto:pravinakalai@gmail.com)

**2.1 Experimental groups**

*Group I – Control group*

Rats given water as vehicle orally, daily for 60 days.

*Group II – Experimental group I (E1)*

Rats given aluminium chloride (100 mg/kg body weight) orally, daily for 60 days.

*Group III – Experimental group II (E2)*

Rats given aluminium chloride 100 mg/kg body weight simultaneously with ginger extract (50mg/kg body weight) orally, daily for 60 days.

*Group IV – Experimental group III (E3)*

Rats given aqueous ginger extract (50mg/kg body weight) orally, daily for 60 days.

During the treatment period, the feed consumed every day was recorded and the body weight was taken on every day and the percent changes were calculated. The animals were treated in the morning. 24 hours after the last treatment, blood samples were collected from the retro-orbital sinus of all animals at the end of the experimental period and EDTA was used as an anti-coagulant. Noncoagulated blood was tested, shortly after collection, for hemoglobin (HB), White blood cells (WBC), Red blood cells (RBC), hematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) using Hematoanalyzer (Siemens).

**3. RESULTS**

Table 1 shows the results of total WBC, RBC, hemoglobin (HB), hematocrit (HCT), and red blood cell indices (MCV, MCH and MCHC) in all groups of rat. The present study reveals that, aluminium chloride administration showed significant elevation in WBC ( $p<0.01$ ) and decline in RBC ( $p<0.01$ ), HB ( $p<0.01$ ), HCT ( $p<0.05$ ) (Fig- 1 and 2). No change was observed in MCV and non significant decrease was noted in MHC and MCHC (Fig-3). On the other hand, administration of ginger extract simultaneously with Al treated animals showed the normal levels similar to control animals. Furthermore, the animal treated with aqueous ginger extract alone does not produce any alteration in all these parameter when compared to control animal.



**Table 1:** Changes in Hematology and RBC'S indices (MCV, MCH, and MCHC) of the aluminium and ginger treated groups compared to control group.

Parameter	Control	AlCl <sub>3</sub>	AlCl <sub>3</sub> , Ginger	Ginger alone
WBC (x10 <sup>9</sup> /L)	6.2 ± 0.326	8.4 ± 0.559 **	5.1 ± 0.664	6.7 ± 0.393
RBC (x10 <sup>12</sup> /L)	5.57 ± 0.383	4.49 ± 0.483 **	5.39 ± 0.178	5.5 ± 0.327
HB (g/dL)	9.08 ± 0.660	7.69 ± 0.753**	8.86 ± 0.308	9.2 ± 0.632
HCT (%)	27.0 ± 1.796	23.23 ± 1.334*	25.84 ± 0.861	27.4 ± 1.704
MCV (fL)	48.6 ± 1.684	46.21 ± 0.736	47.23 ± 0.547	49.0 ± 0.280
MCH (pg)	16.2 ± 0.333	15.6 ± 0.736	16.6 ± 0.231	16.4 ± 0.225
MCHC (g/dL)	33.4 ± 0.254	32.7 ± 0.266	34.2 ± 0.330	33.5 ± 0.369


The results are expressed as Mean ± SEM (n = 6) per treatment and respective control groups.

Levels of significance values are \* $p<0.05$ , \*\* $p<0.01$ , \*\*\* $p<0.001$  compared with control group.

P <0.05 considered to be statistically significant.



**Fig.1 and 2 :** demonstrates the significant changes in the Hematology parameter (WBC, RBC, HB and HCT) of Aluminium administered and ginger treated group compared to control animals.



**Fig. 3 :** Illustrates the significant change in the RBCs indices (MCV, MCHC and MCH) of Aluminium administered and ginger treated group compared to control animals.

**4. DISCUSSION**

Among toxic environmental metals, Al has a remarkable toxic potential for humans. After long term exposure of Al may cause deleterious effect on the morphology and physiology of blood cells. Hematological parameter namely PCV, HB, RBC, WBC and differentials were monitored during the chronic toxicity study in rats because of their role in providing reliable information concerning hematological changes toxicants could cause. The evaluation of hematology and biochemical characteristics has become an important health indicators and means of understanding normal and pathological processes and toxicological impacts<sup>21</sup>. Our study results indicated that Al administration showed significant decline in the concentration of HB, RBC, HCT, etc. It is suggestive that altered peripheral blood composition is a reflection of disrupted hematopoietic process and interfering with different stages of red-cell synthesis and mature red blood cells. However, Al-induced hematological alterations leading to microcytic hypochromic anemia in albino rats on the same blood parameters in rats<sup>22-24</sup>. There was a significant reduction was observed in MCV and RBCs

deformability and significant increments in whole blood viscosity where the animals were treated with 5mg  $Al_2(SO_4)_3$ /kg b.w. for 2 weeks<sup>25</sup>.

The white blood cells are the regulators of the immune system and the increase in WBCs count may be due to generalized immune responses and a protective response to metals stress<sup>26</sup>. In general, Stimulation of lymphopoiesis and/or enhanced release of lymphocytes from lymphomyeloid tissue under toxic stress may lead to an increase in WBCs number<sup>27</sup>. The increased level of WBC in the blood of *C. gariepinus* collected from El-Rahawy drain and this attributed to expose to chronic sewage<sup>28</sup>, and increased WBCs count in fish exposed to copper indicates leucocytosis<sup>29</sup>.

The primary function of erythrocytes is to carry oxygen bound to HB from the lungs to tissues. Furthermore, HB in erythrocytes is an excellent acid-base buffer (for the majority of proteins), thus erythrocytes are mostly responsible for the buffer capacity of whole blood<sup>30</sup>. Significantly decreased HB levels were seen in the Swiss Albino mice after long-term groundwater exposure (180 days) due to heavy metal toxicity<sup>31</sup>. Decreased HB% as well as PCV was reported in wild Libyan Jird, *Meriones libycus* treated with mixture of lead, copper, mercury via drinking water in Riyadh<sup>32</sup>.

Hematocrit (HCT) is used to measure the blood carrying capacity and it is directly associated with the HB% and RBC ability to conduct tissue oxygenation activity. Therefore, HCT shares the reasons for cause of fall in its level with HB% declination in this study. Decrease in HCT may be due to suppresses of bone marrow hematopoietic system, Causing iron deficiency in synthesis of Haem protein of Hemoglobin<sup>33-34</sup>. The low HCT would indicate anaemia or oligohaemia<sup>35</sup>. The changes in the blood HCT value has often been shown to be a good indicator of aluminum toxicity.

Short term exposures to low concentrations of heavy metals mostly induce an increase in the hematological indices<sup>36</sup>. Oliveira Ribeiro *et al.*, observed increase in red blood cells counts in *H. malabaricus* exposed to Methylmercury (MeHg) and attributed this to an increase in the blood oxygen carrying capacity<sup>37</sup>. The RBCs may also be affected in *O. mykiss* after an acute exposure to  $Al^{3+}$ . Further, significant decrease in RBC count and HB were seen in rats and human with high blood lead levels<sup>38-42</sup>.

Significantly decreased RBC, HB levels and HCT values were observed in rats exposed to lead<sup>43-44</sup> and in *Salmo gairdneri*<sup>45</sup>. It was also reported that the decrease in the HCT value of *Salmo gairdneri* blood exposed to cadmium<sup>46</sup>. Moreover, exposure of Zinc to *Oncorhynchus chuskisutsh* caused decrease in HB and HCT values<sup>47</sup>. The reduction in RBCs, HB and HCT might be due to an inhibition of erythropoiesis and hemoglobin synthesis and to an increase in the rate of erythrocytes destruction<sup>48</sup>. Several mechanisms have been proposed for the Al-induced anemia, and the proposed mechanisms appear to involve inhibition of heme synthesis, either by inhibition of enzyme activity or interference with iron incorporation or utilization<sup>49-51</sup>.

Jadhav *et al.*(2007), have reported decreased RBC count after exposing a mixture of heavy metals including (Pb, Cd, Fe, Zn, Cu) in wistar rats<sup>52</sup>. Heavy metal toxicity suppresses the hematopoietic system. Another reason may be decline in erythropoietin synthesis and secretion. In the absence of this hormone, RBC productions were decline considerably<sup>53-54</sup>.

The MCV gives an indication of the status or size of the red blood cells and reflects an abnormal or normal cell division during erythropoiesis. In spite of the increase in the RBC count, a decrease in MCV may show the extent of the shrinking cell size due to Al intoxication<sup>55</sup>. But it was interesting fact in our results that no appreciable changes were seen in MCV in all the experimental groups even though decrease was observed in RBC. The decrease in MCV may indicates that RBC have shrunk, therefore, it is suggestive that the increase in RBC value but the reduced erythrocyte size indicate a high percentage of immature red blood cells in circulation after long term exposure to Al. These can be a marker of anemia with subsequent result of inhibition of erythropoiesis in the hemopoietic system<sup>56</sup>.

It is quite alarming that the use of some medicinal plants seems to pose some health hazards despite their overwhelming health benefits either due to their inherent toxic effect or abuse resulting from over-dosage or prolong usage. Concerns over their safety were raised after incidents of toxicity were reported following consumption of some herbal preparations but the underlying pathogenesis remained cryptic<sup>57-58</sup>.

The rhizome of ginger (*Zingiber officinale*) has been reported to possess a broad-spectrum of prophylactic and therapeutic activities<sup>59</sup>. Ginger is effective in the control of a range of bacterial, viral, fungal and parasitic diseases<sup>60</sup>. It has been extensively studied for the ginger extracts which have a broad range of biological activities; especially antioxidant activities found that ginger significantly lowered lipid peroxidation<sup>61-64</sup>.

Tatfeng and Enitan, [who reported significant increases in total leucocyte count, absolute count of lymphocytes, neutrophils and summation of monocytes, eosinophils and basophils in their work on effects of onion and garlic extracts on some immunologic cells<sup>65</sup>. Yin *et al.*, reported that oral administration of ginger (*Zingiber officinale*) extract increases the phagocytic capability of cells in rainbow trout<sup>66</sup>.

Thus increasing lymphocytes count in the blood following increase in white blood cells count can play an important role in stimulating the immune system of animals. The increase in white blood cells count represents stimulation of the host's immune system. In other words, anti-viral, anti-bacterial and antifungal properties of medicinal plants and symbiotics can be effective in improving immune function and may make the environment unfavorable for invading foreign agents<sup>67</sup>. Herbal immuno-stimulants are substances which activate white blood cells (WBC) and may render fishes more resistant to infectious diseases, by the stimulating phagocytic cells as well as complement lysozyme and antibody responses<sup>68</sup>. Thus, the present study has, however, demonstrated that Al induced anemia can be reversed by the supplementation of ginger extract.

## 5. CONCLUSION

Considering the importance of white blood cells and role of red blood cells in daily human life, the present study was carried out. It was evident in the study that the long term exposure of Al was able to induce the changes in hematological parameters. Any changes made in their structure and number can cause very large physiological changes for animals. On the other hand, supplement administration of ginger extract reversed the Al toxicity effects to the normal level similar to that of control animals. Thus, the present study explores the possibility Al induced anemia can be reversed by the supplementation of ginger extract in rats.

## REFERENCES

1. Foulkes E.C., (1990). Biological Effects of Heavy Metals in Metal Carcinogenesis, Vol. II CRC Press, Florida USA.
2. Jagadeesan G., and Sankarsami Pillai S., (2007). Hepatoprotective effect of taurine against mercury induced toxicity in rat. *J. Environ. Bio.*, 28 753-756.
3. Verstraeten S.V., and Aimo L., (2008). Aluminium and lead: molecular mechanisms of brain toxicity. *Arch. Toxicol.* 82, 789-802.
4. Abbasali K.M., Zhila T., and Farshad N., (2005). Developmental toxicity of aluminium from high doses of  $AlCl_3$  in mice. *J. Applied Res.* 5: 575-579.
5. Cannata, J.J., Briggs, Junor R., and Fell S., (1983). Aluminium hydroxide intake real risk of aluminium toxicity. *Br. Med.* 286: 1937-1938.
6. Lione A., (1985). Aluminium toxicity and the Aluminium containing medication. *J.Pharmacol. Ther.*, 29: 255-285.
7. Kaehny W., Hegg A., and Alfrey A., (1997). Gastrointestinal absorption of aluminium from aluminium containing antacids. *N.Engl.J.Med.*, 296: 1389-1390.
8. Ochmanski W., and Barabasz W., (2000). Aluminium-occurrence and toxicity for organisms. *Przegl. Lek.* 57, 665-668.
9. Exley C., Burgess E., Day J.P., Jeffery E.H., Melethil S. and Yokel R.A. (1996). Aluminium toxicokinetics. *J. Toxicol. Environ. Health.* 48, 569-584.
10. Becaria, A., Campbell, A., and Bondy C.S. Aluminum as a toxicant. *Toxicol. Indust. Health.* 2002: 18:309-320.
11. Farina M., Rotta L.N., Soares F.A., Jardin F., Jacques R., Souza D.O., and Rocha J.B., (2002). Effects of aluminium sulfate on erythropoiesis in rats. *Toxicol. Lett.* 132: 131-139.
12. Masopust J., (2000). Clinical Biochemistry (in Czech). Karolinum, Prague. 832 pp.
13. Kottaimuthu A., Sethupathy S., Manavalan R., and Karar P.K., (2005). Hypolipidemic effect of methahalic extract of

- Dolichos biflorus* (Linn) in high fat diet fed rats. *J. Expl. Biol.* Vol. 43, 522-525.
14. Morakinyo A.O., Adeniyi O.S., Arikawe A.P., (2008). Effects of *Zingiber Officinale* on Reproductive Functions in the Male Rat. *Afric J Biomed Res.* 11(3): 329–34.
  15. Min-Ji, Bak., Seon, O.k., Mira Jun., Woo-Sik Jeong. (2012): 6-shogaol-rich extract from ginger up-regulates the antioxidant defense systems in cells and mice. *Molecules* 17(7):8037–55.
  16. Fisher-Rasmussen W., and Kjaer S., Dahl C., Asping U., (1991). Ginger treatment of Hyperemesis gravidarum. *Eur .J.Obst. Gynecol. Rep. Biol.* 38:19–24.
  17. Sharma G., Nath R., Gill K., (1991). Effects of ethanol on cadmium-induced lipid peroxidation and antioxidant enzymes in rat liver. *Biochem. Pharmacol.* 9–16.
  18. Kamtchoving P., MbongueFndio G., Dimo T., Jatsa H., (2002). Evaluation of androgenic activity of *Zingiberofficinale* and *Pentadiplandrabrazeana* in male rats. *Asian J. Androl.* 4:299–301.
  19. Chrubasik S., Pittler M.H., Roufogalis B.D., (2005). Zingiberis rhizome - A comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine.* 12:684–701.
  20. Khaki A., Fathiazad F., Nouri M., Khaki A., Chelar C., Ozanci, Ghafari-Novin M., Hamadeh M., (2009). The Effects of Ginger on Spermatogenesis and Sperm parameters of Rat. *Iranian J. Reprod. Med.* 7: 7–12.
  21. Saravanan M., Kumar K.P., and. Ramesh M., (2011). Haematological and biochemical responses of freshwater teleost fish *Cyprinus carpio* (Actinopterygii: Cypriniformes) during acute and chronic sublethal exposure to lindane. *Pesticide Biochemistry and Physiology.* 100: 206-211.
  22. Zaman K., Zaman W., Siddique H., (1993). Hematological and enzymatic results of aluminium intoxication in rats. *Comp. Biochem. Physiol.* 150C, 73–76.
  23. Chmielnicka J., Nasiadek M., Lewandowska-Zyndul E., and Pinkowski R., (1996). Effect of aluminum on hematopoiesis after intraperitoneal exposure in rats. *Ecotoxicol. Environ. Saf.* 33(3): 201-6.
  24. Savage D.G., Ogundipe A., Allen R.H., Stabler S.P., and Lindenbaum J., (2000). Etiology and diagnostic evaluation of macrocytosis. *Am J Med Sci.*, 319(6):343 - 352.
  25. Turgut G., B. Kaptanoglu B., S. Turgut S., Y. Enli Y., and O. Genc O., (2004). Effect of chronic aluminum administration on blood and liver iron related parameters in mice. *Yonsei Med. J.*, 45(1): 135-39.
  26. Nussey G., van Vuren J.H.J., and du Preez H.H., (2002). The effect of copper and zinc at neutral and acidic pH on the general haematology and osmoregulation of *Oreochromis mossambicus*. *African Journal of Aquatic Science.* 27(1): 61-84.
  27. El-Sayed Y.S., Saad T.T., and El-Bahr S.M., (2007). Acute intoxication of deltamethrin in monosex Nile tilapia, *Oreochromis niloticus* with special reference to the clinical, biochemical and haematological effects. *Environmental Toxicology and Pharmacology.* 24: 212-217.
  28. Hoeger B., Koellner B., Kotterba G., van den Heuvel M.R., Hitzfeld B., and Dietrich D.R., (2004). Influence of chronic exposure to treated sewage effluent on the distribution of white blood cell populations in rainbow trout (*Oncorhynchus mykiss*) spleen. *Toxicological Sciences,* 82: 97-105.
  29. Dick P., and Dixon D., (1985). Changes in circulating blood cell levels of rainbow trout, *Salmo gairdneri* (Richardson), following acute and chronic exposure to copper. *Journal of Fish Biology,* 26: 475-484.
  30. Ersley A.H., Gabuzda T.G., (1985). Basic Pathophysiology of Blood, 3rd Edition, Philadelphia, WB Saunders Co.
  31. Ravi Kumar Goyal, Kalpana Ojha, Subhasini Sharma, Sharma K.P., (2014). Risk Assessment of Ground water of Sanganer Industrial Area (Jaipur) on Hematology of Swiss Albino Mice. *Journal of Environmental Science, Toxicology and Food Technology* 8(5); 13-17.
  32. Adham K.G., Al-Eisa N.A., Farhood M.H., (2011). Impact of heavy metal pollution on the hemogram and serum biochemistry of the libyan jird, *Meriones libycus*. *Chemosphere.* 84(10):1408–1415.
  33. Klauder D.S., Petering H.G., (1977). Anemia of lead intoxication: A role of copper. *Journal of Nutrition.* 107: 1779-85.
  34. Guyton C., and Hall J.E., (1996). Textbook of Medical Physiology. Ninth edition, Prism Books Pvt Ltd. Bangalore. India. ISBN 81-7286-034-X.
  35. Wepener V., Van Vuren J.H.J., and Du Preez H.H., (1992). The effect of hexavalent chromium at different pH values on the haematology of *Tilapia sparmani* (Chichlidae). *Comparative Biochemistry & Physiology,* C 101, 375–381.
  36. Vutukuru S.S., (2005). Acute Effects of Hexavalent Chromium on Survival, Oxygen Consumption, Hematological Parameters and Some Biochemical Profiles of the Indian Major Carp, *Labeo rohita*. *International Journal of Environmental Research and Public Health,* 2(3): 456-462.
  37. Oliveira Ribeiro C.A., Filipak Neto F., Mela M., Silva P.H., and Randi M.A.F., I.S. Rabitto I.S., Alves Costa J.R.M., and Pelletier E., (2006). Hematological findings in neotropical fish *Hoplias malabaricus* exposed to subchronic and dietary doses of methylmercury, inorganic lead and tributyltin chloride. *Environmental Research,* 101: 74-80.
  38. Allin C.J., and Wilson R.W., (2000). Effects of pre acclimation to aluminium on the physiology and swimming behaviour of juvenile rainbow trout (*Oncorhynchus mykiss*) during a pulsed exposure. *Aquatic Toxicology,* 51: 213-224.
  39. Alexa I.D., Mihalache, I.L., Panaghiu L., and Palade F., (2002). Chronic lead poisoning- a " forgotten" cause of anemia. *Rev. Med. Chir. Soc. Med. Nat. lasi.* 106(4):825-8.
  40. Noori M. M., Heidari Z., Mahmoudzadeh Sagheb H., and Barbarestani M., (2003). Effects of chronic lead acetate intoxication on blood indices of male adult rat. *Daru. Pharm. J.* 11(4): 147-51.
  41. Othman A.I., Al Sharawy S., and El-Missiry M.A., (2004). Role of melatonin in ameliorating lead induced haematotoxicity. *Pharmacol. Res.*, 50(3):301-7.
  42. Toplan S., Ozcelik D., Gulyasar T., and Akyolcu M.C., (2004). Changes in hemorheological parameters due to lead exposure in female rats. *J. Trace. Elem. Med. Biol.*, 18(2):179-82, 2004.
  43. Terayama K., (1993). Effects of lead on electrophoretic mobility, membrane sialic acid, deformability and survival of rat erythrocytes. *Ind Health* 31, 113–126.
  44. Al-Saleh I., (1994). Biochemical and clinical consequences of lead poisoning. *Med Res Rev,* 14: 415–486.
  45. Johansson-Sjoberck M. L., and Larsson A., (1978). The effect of cadmium on the hematology and on the activity of deltaaminolevulinic acid dehydratase (ALA-D) in blood and hematopoietic tissues of the flounder, *Pleuronectes flesus* L. *Environmental Reserch,* 17, 191-204.
  46. Haux C., and Larsson A., (1984). Long-term sublethal physiological effects on Rainbow trout, *Salmo gairdneri*, during exposure to cadmium and after subsequent recovery. *Aquatic Toxicology,* 5, 129-142.
  47. McLeay D.J., (1975). Sensitivity of blood cell counts in juvenil coho salmon (*Oncorhynchus kisutch*) to stressors including sublethal concentrations of pulp mill effluent and zinc. *Journal of the Fisheries Research Board of Canada,* 32, 2357-2364.
  48. Maruyama S., Hirayama C., Yamamoto S., Koda M., Udagawa A., Kadowaki Y., Inoue M., Sagayama A., Umeki K., (2001). Red blood cell status in alcoholic and nonalcoholic liver disease. *J. Lab. Clin. Med.* 138:332-337.
  49. Kaiser L., and Schwartz K.A., 1985. Aluminum-induced anemia. *Am J Kidney Dis.*, 6(5): 348 - 352.
  50. Ganchev T., Dyankov E., Zacharieva R., Pachalieva I., Velikova M., and Kavaldjieva B., (1998). Influence of aluminum on erythropoiesis, iron metabolism and some functional characteristics of erythrocytes in rats. *Acta Physiol Pharmacol Bulg J.*, 23(1): 27- 31.
  51. Han J., Han J., and Dunn M.A., (2000). Effect of dietary aluminum on tissue non-heme iron and ferritin levels in the chick. *Toxicol J.*, 142(2):97-109.
  52. Jadhav S. H., Sarkar S. N., Ram G. C., Tripathi H. C., (2007). Immunosuppressive effect of subchronic exposure to a mixture of eight heavy metals, found as groundwater contaminants in different areas of India, through drinking

- water in male rats. *Archives of environmental contamination and toxicology*. 53(3): 450-458. 2007.
53. Mathur N., Krishnatrey R., Sharma S., Pathak Shipra Sharma K.P., (2003). Certain hematological responses in Swiss albino mice following exposure to textile dye wastewater. *Journal of Environmental Biology*.24(2):161-164.
  54. Sharma S., Sharma S., Sharma K.P., (2006). A comparative study on the toxic effects of textile dye waste waters (Untreated & Treated) on mortality and RBC of a freshwater fish *Gambusia affinis* (Baird & Gerard). *Journal of Environmental Biology*. 27(4): 623-628.
  55. Alwan S.F., Hadi A. A., and Shokr A. E., (2009). Alterations in haematological parameter of fresh water fish *Tilapia Zilli* exposed to Aluminium. *Journal of Science and Its Applications*, 3(1):12-19.
  56. Lavanya S., Ramesh M., Kavitha C., and Malarvizhi A., (2011). Hematological, biochemical and ionoregulatory responses of Indian major carp *Catla catla* during chronic sublethal exposure to inorganic arsenic. *Chemosphere*, 82, 977–985.
  57. Favreau J.T., Ryu M.L., Braunstein G., Orshansky G., Park S.S., Coody G.L., (2002). Severe hepatotoxicity associated with the dietary supplement LipoKinetix. *Ann Intern Med*. 136:590–595.
  58. Stickel F., Patsenker, E., and Schuppan D., (2005). Herbal hepatotoxicity. *J. Hepatol*. 43: 901-910.
  59. Ernst E., and Pittler M. H., (2000). Efficacy of ginger for nausea and vomiting: A systematic review of randomized clinical trials. *Br. J. Anaesth*. 84:367-371.
  60. Endo K., Kanno E., and Oshima Y., (1990). Structures of antifungal diarylheptenons, gingerones A, B, C and isogingerenones, isolated from the rhizomes of *Zingiber officinale*, *Phytochemistry* 29:797-799.
  61. Miller J.K., Brzezinska-Slebodzinska E., Madsen F.C., (1993). Oxidative stress, antioxidant and animals function. *J. Dairy Sci*. 76: 2812–2823.
  62. Ahmed R.S., Seth V., and Banerjee, B.D., (2000). Influence of dietary ginger on antioxidant defence system in rat: comparison with ascorbic acid. *Ind. J. Exp. Biol*, 38:604-606.
  63. Agrawal M., Walia S., Dhingra S., and Khambay B.P.S., (2001). Insect Growth inhibition antifeedant and antifungal activity of compounds isolated derived from *Zingiber officinale* Roscoe, ginger rhizome. *Pest Manag. Sci*. 57: 289-300.
  64. Martins A.P., Salgueiro L., Goncalves M.J., Cunchada A.P., and Vila R., (2001). Essential oil composition and antimicrobial activity of three *Zingiberaceae* from S. Tome Principe. *Planta Medicine*; 67:580-584.
  65. Tاتفeng Y.M., Enitan S.S., (2012). Effects of onion and garlic extracts on some immunologic cells. *African Journal of Traditional, Complementary and Alternative medicine*, 9(3); 374-379.
  66. Yin G., Jeney G., Racz T., XU. and Jun X., (2006). Effect of two Chinese herbs Astragalusradix and Scutellaria radix) on non specific immune response of tilapia, *Oreochromis niloticus*, *Aquaculture* 23,P,39-47.
  67. Al-Kassi G.A.M., (2009). Influence of plant exteracts derived from thyme and cinnamon on broiler performance. *Pak. Vet. J*. 29(4): 169-173.
  68. Secombes C.J., and Olivier G., (1997). Furunculosis. Academic Press, New York, pp. 269-296.